

# Notes on Matrix Valued Paraproducts

Tao MEI <sup>1</sup>

**Abstract** Denote by  $M_n$  the algebra of  $n \times n$  matrices. We consider the dyadic paraproducts  $\pi_b$  associated with  $M_n$  valued functions  $b$ , and show that the  $L^\infty(M_n)$  norm of  $b$  does not dominate  $\|\pi_b\|_{L^2(\ell_n^2) \rightarrow L^2(\ell_n^2)}$  uniformly over  $n$ . We also consider paraproducts associated with noncommutative martingales and prove that their boundedness on bounded noncommutative  $L^p$ -martingale spaces implies their boundedness on bounded noncommutative  $L^q$ -martingale spaces for all  $1 < p < q < \infty$ .

## 1 Introduction

Denote by  $M_n$  the algebra of  $n \times n$  matrices. Let  $(\mathbb{T}, \mathcal{F}_k, dt)$  be the unit circle with Haar measure and the usual dyadic filtration. Let  $b$  be an  $M_n$  valued function on  $\mathbb{T}$ . The matrix valued dyadic paraproduct associated with  $b$ , denoted by  $\pi_b$ , is the operator defined as

$$\pi_b(f) = \sum_k (d_k b)(E_{k-1} f), \quad \forall f \in L^2(\ell_n^2). \quad (1.1)$$

Here  $E_k f$  is the conditional expectation of  $f$  with respect to  $\mathcal{F}_k$ , i.e. the unique  $\mathcal{F}_k$ -measurable function such that

$$\int_F E_k f dt = \int_F f dt, \quad \forall F \in \mathcal{F}_k.$$

And  $d_k b$  is defined to be  $E_k b - E_{k-1} b$ .

In the classical case (when  $b$  is a scalar valued function), paraproducts are usually considered as dyadic singular integrals and play important roles in the proof of the classical T(1) theorem. It is well known that

$$\|\pi_b\|_{L^2 \rightarrow L^2} \simeq \|b\|_{BMO_d},$$

where  $BMO_d$  denotes the dyadic BMO norm defined as

$$\|b\|_{BMO_d} = \sup_m \|E_m \sum_{k=m}^{\infty} |d_k b|^2\|_{L^\infty}^{\frac{1}{2}}.$$

And by the Calderón-Zygmund decomposition and the Marcinkiewicz interpolation theorem, we have  $\|\pi_b\|_{L^p \rightarrow L^p} \simeq \|\pi_b\|_{L^p \rightarrow L^p} \simeq \|b\|_{BMO_d}$  for all  $1 < p < \infty$ .

When  $b$  is  $M_n$  valued, it was proved by Katz ([4]) and independently by Nazarov, Treil and Volberg ([8], see [10] for another proof by Pisier) that

$$\|\pi_b\|_{L^2(\ell_n^2) \rightarrow L^2(\ell_n^2)} \leq c \log(n+1) \|b\|_{BMO_c}. \quad (1.2)$$

---

<sup>1</sup>Partially supported by the National Science Foundation 0200690.

Here  $\|\cdot\|_{\text{BMO}_c}$  is the column BMO norm defined by

$$\|b\|_{\text{BMO}_c} = \sup_m \left\| E_m \sum_{k=m}^{\infty} (d_k b)^* (d_k b) \right\|_{L^\infty(M_n)}^{\frac{1}{2}},$$

where  $(d_k b)^*$  is the adjoint of  $d_k b$ . Nazarov, Pisier, Treil and Volberg ([7]) proved later that the constant  $c \log(n+1)$  in (1.2) is optimal. Thus the  $\text{BMO}_c$  norm does not dominate  $\|\pi_b\|_{L^2(\ell_n^2) \rightarrow L^2(\ell_n^2)}$  uniformly over  $n$ .

Can we expect something weaker? In particular, does there exist a constant  $c$  independent of  $n$  such that, for every  $n \in \mathbb{N}$ ,

$$\|\pi_b\|_{L^2(\ell_n^2) \rightarrow L^2(\ell_n^2)} \leq c \|b\|_{L^\infty(M_n)}? \quad (1.3)$$

Some known facts made (1.3) look hopeful. For example, the Hankel operator associated with the  $M_n$  valued function  $b$  has a norm equivalent to  $\|b\|_{(H^1(S^1))^*}$ . Here  $\|\cdot\|_{(H^1(S^1))^*}$  denotes the dual norm on the trace class valued Hardy space  $H^1(S^1)$ . And S. Petermichl proved a close relation between  $\pi_b$  and the Hankel operators associated with  $b$  (see [9]).

In this paper, we prove the following theorem, which shows there does not exist any constant  $c$  independent of  $n$  such that (1.3) holds.

**Theorem 1.1** *For every  $n \in \mathbb{N}$ , there exists an  $M_n$  valued function  $b$  with  $\|b\|_{L^\infty(M_n)} \leq 1$  but such that*

$$\|\pi_b\|_{L^2(\ell_n^2) \rightarrow L^2(\ell_n^2)} \geq c \log(n+1),$$

where  $c > 0$  is independent of  $n$ .

This also gives a new proof that the constant  $c \log(n+1)$  in (1.2) is optimal.

Denote by  $S^p$  the Schatten  $p$  class on  $\ell^2$ . For  $f \in L^p(S^p)$ , we define  $\pi_b(f)$  as in (1.1) also. As pointed out in [10], it is easy to check that  $\|\pi_b\|_{L^2(S^2) \rightarrow L^2(S^2)} = \|\pi_b\|_{L^2(\ell^2) \rightarrow L^2(\ell^2)}$ . For scalar valued  $b$ , as we mentioned previously, we have  $\|\pi_b\|_{L^p \rightarrow L^p} \simeq \|\pi_b\|_{L^q \rightarrow L^q}$ . We wonder if this is still true for matrix valued  $b$ , i.e. if  $\pi_b$ 's boundedness on  $L^p(S^p)$  implies their boundedness on  $L^q(S^q)$  for all  $1 < p, q < \infty$ .

More generally, we can consider paraproducts associated with noncommutative martingales. Let  $\mathcal{M}$  be a finite von Neumann algebra with a normalized faithful trace  $\tau$ . For  $1 \leq p < \infty$ , we denote by  $L^p(\mathcal{M})$  the noncommutative  $L^p$  space associated with  $(\mathcal{M}, \tau)$ . Recall the norm in  $L^p(\mathcal{M})$  is defined as

$$\|f\|_p = (\tau|f|^p)^{\frac{1}{p}}, \quad \forall f \in L^p(\mathcal{M}),$$

where  $|f| = (f^* f)^{\frac{1}{2}}$ . For convenience, we usually set  $L^\infty(\mathcal{M}) = \mathcal{M}$  equipped with the operator norm  $\|\cdot\|_{\mathcal{M}}$ . Let  $\mathcal{M}_k$  be an increasing filtration of von Neumann subalgebras of  $\mathcal{M}$  such that  $\cup_{k \geq 0} \mathcal{M}_k$  generates  $\mathcal{M}$  in the  $w^*$ -topology. Denote by  $E_k$  the conditional expectation of  $\mathcal{M}$  with respect to  $\mathcal{M}_k$ .  $E_k$  is a norm 1 projection of  $L^p(\mathcal{M})$  onto

$L^p(\mathcal{M}_k)$ . For  $1 \leq p \leq \infty$ , a sequence  $f = (f_k)_{k \geq 0}$  with  $f_k \in L^p(\mathcal{M}_k)$  is called a bounded noncommutative  $L^p$ -martingale, denoted by  $(f_k)_{k \geq 0} \in L^p(\mathcal{M})$ , if  $E_k f_m = f_k, \forall k \leq m$  and

$$\|(f_k)_{k \geq 0}\|_{L^p(\mathcal{M})} = \sup_k \|f_k\|_{L^p(\mathcal{M})} < \infty.$$

Because of the uniform convexity of the space  $L^p(\mathcal{M})$ , for  $1 < p < \infty$ , we can and will identify the space of all bounded  $L^p(\mathcal{M})$ -martingales with  $L^p(\mathcal{M})$  itself. In particular, for any  $f \in L^p(\mathcal{M})$ , set  $f_k = E_k f$ , then  $f = (f_k)_{k \geq 0}$  is a bounded  $L^p(\mathcal{M})$ -martingale and  $\|(f_k)_{k \geq 0}\|_{L^p(\mathcal{M})} = \|f\|_{L^p(\mathcal{M})}$ . Denote by  $d_k f = E_k f - E_{k-1} f$ .

We say an increasing filtration  $\mathcal{M}_k$  is “regular” if there exists a constant  $c > 0$  such that, for any  $m, a \in \mathcal{M}_m, a \geq 0$ ,

$$\|a\|_\infty \leq c \|E_{m-1} a\|_\infty.$$

For  $\mathcal{M}$  with a regular filtration  $\mathcal{M}_k$ ,  $b \in L^2(\mathcal{M})$ , we define paraproducts  $\pi_b, \tilde{\pi}_b$  as operators for bounded  $L^p(\mathcal{M})$  ( $1 < p < \infty$ )-martingales  $f = (f_k)_{k \geq 0}$  as

$$\pi_b(f) = \sum_k d_k b f_{k-1}, \quad \tilde{\pi}_b(f) = \sum_k f_{k-1} d_k b.$$

We prove the following result for  $\pi_b$  and  $\tilde{\pi}_b$  :

**Theorem 1.2** *Let  $1 < p < q < \infty$ , if  $\tilde{\pi}_b$  and  $\pi_b$  are both bounded on  $L^p(\mathcal{M})$  then they are both bounded on  $L^q(\mathcal{M})$ .*

We still do not know what happens when  $p > q$ .

## 2 Proof of Theorem 1.1 and Application to “Sweep” functions.

Denote by  $tr$  the usual trace on  $M_n$  and  $S_n^p$  ( $1 \leq p < \infty$ ) the Schatten  $p$  classes on  $\ell_n^2$ .

**Proof of Theorem 1.1.** Let  $c(n)$  be the best constant such that

$$\|\pi_b\|_{L^2(\ell_n^2) \rightarrow L^2(\ell_n^2)} \leq c(n) \|b\|_{L^\infty(M_n)}, \quad \forall b \in L^\infty(M_n).$$

Denote by  $T$  the triangle projection on  $S_n^1$ , we are going to show

$$\|T\|_{S_n^1 \rightarrow S_n^1} \leq c(n).$$

Once this is proved, we are done since  $\|T\|_{S_n^1 \rightarrow S_n^1} \sim \log(n+1)$  (see [5]). Note that every  $A$  in the unit ball of  $S_n^1$  can be written as

$$A = \sum_m \lambda^{(m)} \alpha^{(m)} \otimes \beta^{(m)}$$

with  $\sum_m \lambda^{(m)} \leq 1$ ,  $\sup_m \{\|\alpha^{(m)}\|_{\ell_n^2}, \|\beta^{(m)}\|_{\ell_n^2}\} \leq 1$ . Therefore, we only need to show

$$\|T(\alpha \otimes \beta)\|_{S_n^1} \leq c(n) \|\alpha\|_{\ell_n^2} \|\beta\|_{\ell_n^2}, \quad \forall \alpha = (\alpha_k)_k, \beta = (\beta_k)_k \in \ell_n^2. \quad (2.4)$$

Let  $D$  be the diagonal  $M_n$  valued function defined as

$$D = \sum_{i=1}^n r_i e_i \otimes e_i$$

where  $r_i$  is the  $i$ -th Rademacher function on  $\mathbb{T}$  and  $(e_i)_{i=1}^n$  is the canonical basis of  $\ell_n^2$ . Given  $\alpha = (\alpha_k)_k, \beta = (\beta_k)_k \in \ell_n^2$ , let

$$f = D\alpha, g = D\beta.$$

Then  $f, g \in L^2(\ell_n^2)$ , and

$$\|f\|_{L^2(\ell_n^2)} = \|\alpha\|_{\ell_n^2}, \|g\|_{L^2(\ell_n^2)} = \|\beta\|_{\ell_n^2}. \quad (2.5)$$

It is easy to verify

$$\sum_k E_{k-1} f \otimes d_k g = D \left( \sum_{i < j \leq n} \alpha_i \beta_j e_i \otimes e_j \right) D.$$

and

$$\left\| \sum_k E_{k-1} f \otimes d_k g \right\|_{L^1(S_n^1)} = \left\| \sum_{i < j \leq n} \alpha_i \beta_j e_i \otimes e_j \right\|_{S_n^1} = \|T(\alpha \otimes \beta)\|_{S_n^1}. \quad (2.6)$$

On the other hand, by duality between  $L^1(S_n^1)$  and  $L^\infty(M_n)$ , we have,

$$\begin{aligned} \left\| \sum_k E_{k-1} f \otimes d_k g \right\|_{L^1(S_n^1)} &= \sup \left\{ \operatorname{tr} \int \sum_k d_k b (E_{k-1} f \otimes d_k g), \|b\|_{L^\infty(M_n)} \leq 1 \right\} \\ &\leq \sup \left\{ \|\pi_b(f)\|_{L^2(\ell_n^2)} \|g\|_{L^2(\ell_n^2)}, \|b\|_{L^\infty(M_n)} \leq 1 \right\} \\ &\leq c(n) \|f\|_{L^2(\ell_n^2)} \|g\|_{L^2(\ell_n^2)}. \end{aligned} \quad (2.7)$$

Combining (2.7), (2.5) and (2.6) we get (2.4) and the proof is complete. ■

Recall that the square function of  $b$  is defined as

$$S(b) = \left( \sum_k |d_k b|^2 \right)^{\frac{1}{2}}.$$

The so called “sweep” function is just the square of the square function, for this reason we denote it by  $S^2(b)$ ,

$$S^2(b) = \sum_k |d_k b|^2.$$

In the classical case, we know that

$$\|S(b)\|_{BMO_d} \leq c\|b\|_{BMO_d} \quad (2.8)$$

$$\|S^2(b)\|_{BMO_d} \leq c\|b\|_{BMO_d}^2 \quad (2.9)$$

When considering square functions  $S(b)$  for  $M_n$  valued functions  $b$ , a similar result remains true with an absolute constant.

**Proposition 2.3** *For any  $n \in \mathbb{N}$ , and any  $M_n$  valued function  $b$ , we have*

$$\|S(b)\|_{BMO_c} \leq \sqrt{2}\|b\|_{BMO_c}$$

**Proof.** Since we are in the dyadic case, we have

$$\begin{aligned} \|S(b)\|_{BMO_c}^2 &\leq 2 \sup_m \|E_m[(S(b) - E_m S(b))^*(S(b) - E_m S(b))]\|_{L^\infty(M_n)} \\ &= 2 \sup_m \|E_m S^2(b) - (E_m S(b))^2\|_{L^\infty(M_n)} \end{aligned}$$

Note

$$E_m S^2(b) - \sum_{k=1}^m |d_k b|^2 \geq E_m S^2(b) - (E_m S(b))^2 \geq 0.$$

We get

$$\begin{aligned} \|S(b)\|_{BMO_c}^2 &\leq 2 \sup_m \|E_m S^2(b) - \sum_{k=1}^m |d_k b|^2\|_{L^\infty(M_n)} \\ &= 2 \sup_m \|E_m \sum_{k=m+1}^\infty |d_k b|^2\|_{L^\infty(M_n)} \\ &\leq 2\|b\|_{BMO_c}^2. \blacksquare \end{aligned}$$

Matrix valued sweep functions have been studied in [1], [2] etc. Unlike in the case of square functions, it is proved in [1] that the best constant  $c_n$  such that

$$\|S^2(b)\|_{BMO_c} \leq c_n \|b\|_{BMO_c}^2 \quad (2.10)$$

is  $c \log(n+1)$ . The following result shows that the best constant  $c_n$  is still  $c \log(n+1)$  even if we replace  $\|\cdot\|_{BMO_c}$  by the bigger norm  $\|\cdot\|_{L^\infty(M_n)}$  in the right side of (2.10).

**Theorem 2.4** *For every  $n \in \mathbb{N}$ , there exists an  $M_n$  valued function  $b$  with  $\|b\|_{L^\infty(M_n)} \leq 1$  but such that*

$$\|S^2(b)\|_{BMO_c} \geq c \log(n+1).$$

**Proof.** Consider a function  $b$  that works for the statement of Theorem 1.1. Then  $\|b\|_{L^\infty(M_n)} \leq 1$  and there exists a function  $f \in L^2(S_n^2)$ , such that  $\|f\|_{L^2(S_n^2)} \leq 1$  and

$$\left\| \sum_k d_k b E_{k-1} f \right\|_{L^2(S_n^2)} \geq c \log(n+1). \quad (2.11)$$

We compute the square of the left side of (2.11) and get

$$\begin{aligned} & \left\| \sum_k d_k b E_{k-1} f \right\|_{L^2(S_n^2)}^2 \\ &= \text{tr} \int \sum_k |d_k b|^2 E_{k-1} f E_{k-1} f^* \\ &= \text{tr} \int \sum_k |d_k b|^2 \left( \sum_{i < k} |d_i f^*|^2 + \sum_{i < k} E_{i-1} f d_i f^* + \sum_{i < k} d_i f E_{i-1} f^* \right) \\ &= \text{tr} \int \sum_i \left( \sum_{k > i} |d_k b|^2 \right) |d_i f^*|^2 + \text{tr} \int \sum_i \left( \sum_{k > i} |d_k b|^2 \right) (E_{i-1} f d_i f^* + d_i f E_{i-1} f^*) \\ &= I + II \end{aligned}$$

For  $I$ , note  $|d_i f^*|^2$  is  $\mathcal{F}_i$  measurable, we have

$$\begin{aligned} I &= \text{tr} \int \sum_i E_i \left( \sum_{k > i} |d_k b|^2 \right) |d_i f^*|^2 \\ &\leq \sup_i \|E_i \left( \sum_{k > i} |d_k b|^2 \right)\|_{L^\infty(M_n)} \left( \text{tr} \int \sum_i |d_i f^*|^2 \right) \\ &\leq \|b\|_{BMO_c}^2 \|f\|_{L^2(S_n^2)}^2 \leq 4 \end{aligned}$$

For  $II$ , note  $E_{i-1} f d_i f^* + d_i f E_{i-1} f^*$  is a martingale difference and  $\sum_{k \leq i} |d_k|^2$  is  $\mathcal{F}_{i-1}$  measurable since we are in the dyadic case, we get

$$\begin{aligned} II &= \text{tr} \int \sum_i S^2(b) (E_{i-1} f d_i f^* + d_i f E_{i-1} f^*) \\ &= \text{tr} \int \sum_i d_i (S^2(b)) (E_{i-1} f d_i f^* + d_i f E_{i-1} f^*) \\ &\leq 2 \left\| \sum_i d_i (S^2(b)) E_{i-1} f \right\|_{L^2(S_n^2)} \|f\|_{L^2(S_n^2)} \\ &\leq 2 \|\pi_{S^2(b)}\|_{L^2(S_n^2) \rightarrow L^2(S_n^2)} \\ &\leq 2c \log(n+1) \|S^2(b)\|_{BMO_c}. \end{aligned}$$

We used (1.2) in the last step. Combining this with (2.11), we get

$$c \log(n+1) \leq \left\| \sum_k d_k b E_{k-1} f \right\|_{L^2(S_n^2)}^2 \leq 4 + 2c \log(n+1) \|S^2(b)\|_{BMO_c}$$

Thus

$$\|S^2(b)\|_{\text{BMO}_c} \geq c \log(n+1).$$

This completes the proof. ■

### 3 Proof of Theorem 1.2.

We keep the notations introduced in the end of Section 1. Recall BMO spaces of noncommutative martingales are defined for  $x = (x_k) \in L^2(\mathcal{M})$  as below (see [?], [?]):

$$\begin{aligned} \text{BMO}_c(\mathcal{M}) &= \{x : \|x\|_{\text{BMO}_c(\mathcal{M})} = \sup_n \left\| E_n \left| \sum_{k=n}^{\infty} d_k x \right|^2 \right\|_{\mathcal{M}}^{\frac{1}{2}} < \infty\}; \\ \text{BMO}_r(\mathcal{M}) &= \{x : \|x\|_{\text{BMO}_r(\mathcal{M})} = \|x^*\|_{\text{BMO}_c(\mathcal{M})} < \infty\}; \\ \text{BMO}_{cr}(\mathcal{M}) &= \{x : \|x\|_{\text{BMO}_{cr}(\mathcal{M})} = \max\{\|x\|_{\text{BMO}_c(\mathcal{M})}, \|x\|_{\text{BMO}_r(\mathcal{M})}\} < \infty\}. \end{aligned}$$

When  $\mathcal{M} = L^\infty(M_n)$ ,  $\text{BMO}_c(\mathcal{M})$  is just  $\text{BMO}_c$  considered in Section 1 and 2. In this section, for noncommutative martingale  $b$ , we consider  $\pi_b$  and  $\tilde{\pi}_b$  as operators on bounded noncommutative  $L^p$ -martingale spaces introduced in Section 1. We will need the following interpolation result and the John-Nirenberg theorem for noncommutative martingales proved by Junge and Musat recently (see [3], [6]).

**Theorem 3.5** (*Musat*) For  $1 \leq p \leq q < \infty$ ,

$$(\text{BMO}_{cr}(\mathcal{M}), L_p(\mathcal{M}))_\theta = L_q(\mathcal{M}), \text{ with } \theta = \frac{p}{q}.$$

**Theorem 3.6** (*Junge, Musat*) For any  $1 \leq q < \infty$  and any  $g = (g_k)_k \in \text{BMO}_{cr}(\mathcal{M})$ , there exist  $c_q, c'_q > 0$  such that

$$c'_q \|g\|_{\text{BMO}_{cr}} \leq \sup_{m \in \mathbb{N}} \sup_{a \in \mathcal{M}_m, \tau(|a|^q) \leq 1} \left\{ \left\| \sum_{k \geq m} d_k g a \right\|_{L^q(\mathcal{M})}, \left\| \sum_{k \geq m} a d_k g \right\|_{L^q(\mathcal{M})} \right\} \leq c_q \|g\|_{\text{BMO}_{cr}}. \quad (3.12)$$

In fact, the formula above is proved for  $q \geq 2$  in [3]. It is not hard to show that it is also true for  $1 \leq q < 2$ . In the following, we give a simpler proof of it in the tracial case.

**Proof.** Note for any  $g \in \text{BMO}_{cr}(\mathcal{M})$ ,

$$\|g\|_{\text{BMO}_{cr}(\mathcal{M})} = \sup_{m \in \mathbb{N}} \sup_{a \in \mathcal{M}_m, \tau(|a|^2) \leq 1} \left\{ \left\| \sum_{k \geq m} d_k g a \right\|_{L^2(\mathcal{M})}, \left\| \sum_{k \geq m} a d_k g \right\|_{L^2(\mathcal{M})} \right\}.$$

We get  $c_2 = c'_2 = 1$ . Note for  $p, r, s$  with  $1/p = 1/r + 1/s$  and  $a \in L^p(\mathcal{M})$ ,  $\|a\|_{L^p(\mathcal{M})} \leq 1$ , there exist  $b, c$  such that  $a = bc$  and  $\|b\|_{L^r(\mathcal{M})} \leq 1$ ,  $\|c\|_{L^s(\mathcal{M})} \leq 1$ . By Hölder's inequality we then get  $c_q = 1$  for  $1 \leq q < 2$  and  $c'_q = 1$  for  $2 < q < \infty$ . Thus for  $2 < q < \infty$ , we

only need to prove the second inequality of (3.12). And, for  $1 \leq q < 2$ , we only need to prove the first inequality of (3.12). Fix  $g \in BMO_{cr}(\mathcal{M})$ ,  $m \in \mathbb{N}$ , consider the left multiplier  $L_m$  and the right multiplier  $R_m$  defined as

$$L_m(a) = \sum_{k \geq m} d_k g a \text{ and } R_m(a) = \sum_{k \geq m} a d_k g, \quad \forall a \in \mathcal{M}_m.$$

It is easy to check that

$$\begin{aligned} \sup_m \|L_m\|_{L^2(\mathcal{M}_m) \rightarrow L^2(\mathcal{M})} &= \|g\|_{BMO_c}, \\ \sup_m \|L_m\|_{L^\infty(\mathcal{M}_m) \rightarrow BMO_{cr}} &\leq \|g\|_{BMO_{cr}}; \\ \sup_m \|R_m\|_{L^2(\mathcal{M}_m) \rightarrow L^2(\mathcal{M})} &= \|g\|_{BMO_r}, \\ \sup_m \|R_m\|_{L^\infty(\mathcal{M}_m) \rightarrow BMO_{cr}} &\leq \|g\|_{BMO_{cr}}. \end{aligned}$$

Thus  $L_m, R_m$  extend to bounded operators from  $L^2(\mathcal{M}_m)$  to  $L^2(\mathcal{M})$ , as well as from  $L^\infty(\mathcal{M}_m)$  to  $BMO_{cr}(\mathcal{M})$ . By Musat's interpolation result Theorem 3.5, we get  $L_m$  and  $R_m$  are bounded from  $L^q(\mathcal{M}_m)$  to  $L^q(\mathcal{M})$  and their operator norms are smaller than  $c_q \|g\|_{BMO_{cr}}$ , for all  $2 \leq q < \infty$ . By taking supremum over  $m$ , we prove the second inequality of (3.12) for  $q \geq 2$ .

For  $1 \leq q < 2$ , by interpolation again, for  $\theta = \frac{q}{2}$  and some  $c_q'' > 0$ ,

$$\begin{aligned} \|L_m\|_{L^2(\mathcal{M}_m) \rightarrow L^2(\mathcal{M})} &\leq c_q'' \|L_m\|_{L^q(\mathcal{M}_m) \rightarrow L^q(\mathcal{M})}^\theta \|L_m\|_{L^\infty(\mathcal{M}_m) \rightarrow BMO_{cr}}^{1-\theta} \\ &\leq c_q'' \|L_m\|_{L^q(\mathcal{M}_m) \rightarrow L^q(\mathcal{M})}^\theta \|g\|_{BMO_{cr}}^{1-\theta}, \\ \|R_m\|_{L^2(\mathcal{M}_m) \rightarrow L^2(\mathcal{M})} &\leq c_q'' \|R_m\|_{L^q(\mathcal{M}_m) \rightarrow L^q(\mathcal{M})}^\theta \|R_m\|_{L^\infty(\mathcal{M}_m) \rightarrow BMO_{cr}}^{1-\theta} \\ &\leq c_q'' \|R_m\|_{L^q(\mathcal{M}_m) \rightarrow L^q(\mathcal{M})}^\theta \|g\|_{BMO_{cr}}^{1-\theta}. \end{aligned}$$

Thus

$$\begin{aligned} \|g\|_{BMO_{cr}} &= \max\{\sup_m \|L_m\|_{L^2(\mathcal{M}_m) \rightarrow L^2(\mathcal{M})}, \sup_m \|R_m\|_{L^2(\mathcal{M}_m) \rightarrow L^2(\mathcal{M})}\} \\ &\leq c_q'' \|g\|_{BMO_{cr}}^{1-\theta} \sup_m \{\|L_m\|_{L^q(\mathcal{M}_m) \rightarrow L^q(\mathcal{M})}^\theta, \|R_m\|_{L^q(\mathcal{M}_m) \rightarrow L^q(\mathcal{M})}^\theta\}. \end{aligned}$$

This gives the first inequality of (3.12) with  $c'_q = (c_q'')^{-\frac{1}{\theta}}$  for  $1 \leq q < 2$ . ■

Recall that we say a filtration  $\mathcal{M}_k$  is “regular” if, for some  $c > 0$ ,  $\|a\|_\infty \leq c \|E_{m-a}\|_\infty$ ,  $\forall m \in \mathbb{N}, a \geq 0, a \in \mathcal{M}_m$ .

**Lemma 3.7** *For any regular filtration  $\mathcal{M}_k$ , we have*

$$\|b\|_{BMO_{cr}(\mathcal{M})} \leq c_p \max\{\|\pi_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})}, \|\tilde{\pi}_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})}\}, \quad \forall 1 \leq p < \infty. \quad (3.13)$$

**Proof.** Note, for any  $b \in BMO_{cr}(\mathcal{M})$  with respect to the regular filtration  $\mathcal{M}_k$ ,

$$\|b\|_{BMO_{cr}(\mathcal{M})} \leq c \sup_{m \in \mathbb{N}} \sup_{\tau a^2 \leq 1, a \in \mathcal{M}_m} \left\{ \left\| \sum_{k > m} d_k b a \right\|_{L^2(\mathcal{M})}, \left\| \sum_{k > m} a d_k b \right\|_{L^2(\mathcal{M})} \right\}.$$



Similar to the proof of Theorem 3.6, we can get,

$$c'_q \|b\|_{BMO_{cr}} \leq \sup_{m \in \mathbb{N}} \sup_{a \in \mathcal{M}_m, \tau|a|^q \leq 1} \{ \left\| \sum_{k>m} d_k b a \right\|_{L^q(\mathcal{M})}, \left\| \sum_{k>m} a d_k b \right\|_{L^q(\mathcal{M})} \} \leq c_q \|b\|_{BMO_{cr}}. \quad (3.14)$$

On the other hand, by considering  $\pi_b(a), \tilde{\pi}_b(a)$  for  $a \in \mathcal{M}_m$ ,  $\|a\|_{L^p(\mathcal{M})} \leq 1$ , we have

$$\begin{aligned} & \sup_{a \in \mathcal{M}_m, \tau|a|^q \leq 1} \{ \left\| \sum_{k>m} d_k b a \right\|_{L^p(\mathcal{M})}, \left\| \sum_{k>m} a d_k b \right\|_{L^p(\mathcal{M})} \} \\ & \leq 2 \max \{ \|\pi_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})}, \|\tilde{\pi}_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})} \}. \end{aligned}$$

Taking supremum over  $m$  in the inequality above, we get (3.13) by (3.14). ■

**Lemma 3.8** For  $1 < p < \infty$ , we have

$$\|\pi_b\|_{L^\infty(\mathcal{M}) \rightarrow BMO_{cr}(\mathcal{M})} \leq c_p (\|\pi_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})} + \|b\|_{BMO_r(\mathcal{M})}). \quad (3.15)$$

$$\|\tilde{\pi}_b\|_{L^\infty(\mathcal{M}) \rightarrow BMO_{cr}(\mathcal{M})} \leq c_p (\|\tilde{\pi}_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})} + \|b\|_{BMO_c(\mathcal{M})}). \quad (3.16)$$

**Proof.** We prove (3.15) only. Fix a  $f \in L^\infty(\mathcal{M})$  with  $\|f\|_{L^\infty(\mathcal{M})} \leq 1$ . We have

$$\begin{aligned} & \left\| E_m \sum_{k \geq m} |d_k b E_{k-1} f|^2 \right\|_{L^\infty(\mathcal{M})} \\ &= \sup \{ \tau E_m \sum_{k \geq m} |d_k b E_{k-1} f|^2 a, a \in \mathcal{M}_m, a \geq 0, \tau a \leq 1 \} \\ &= \sup \{ \tau \sum_{k \geq m} (d_k b E_{k-1} f a^{\frac{1}{p}})^* (d_k b E_{k-1} f a^{\frac{1}{q}}), a \in \mathcal{M}_m, a \geq 0, \tau a \leq 1 \} \\ &\leq \sup_a \left\| d_m b E_{m-1} f a^{\frac{1}{p}} + \sum_{k>m} d_k b E_{k-1} (f a^{\frac{1}{p}}) \right\|_{L^p(\mathcal{M})} \left\| \sum_{k \geq m} d_k b E_{k-1} f a^{\frac{1}{q}} \right\|_{L^q(\mathcal{M})} \end{aligned}$$

Note  $\|d_m b E_{m-1} f a^{\frac{1}{p}}\|_{L^p(\mathcal{M})} \leq \|d_m b\|_{\mathcal{M}} \leq \|b\|_{BMO_r}$ . By (3.12) we get

$$\left\| E_m \sum_{k \geq m} |d_k b E_{k-1} f|^2 \right\|_{L^\infty(\mathcal{M})} \leq c_q (\|b\|_{BMO_r} + \|\pi_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})}) \|\pi_b(f)\|_{BMO_{cr}(\mathcal{M})} \quad (3.17)$$

Taking supremum over  $m$  in (3.17), we get

$$\|\pi_b(f)\|_{BMO_c(\mathcal{M})}^2 \leq c_q (\|b\|_{BMO_r} + \|\pi_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})}) \|\pi_b(f)\|_{BMO_{cr}(\mathcal{M})}.$$

On the other hand, since  $(E_{m-1}f)(E_{m-1}f)^* \leq 1$ , we have

$$\|\pi_b(f)\|_{BMO_r(\mathcal{M})} \leq \|b\|_{BMO_r(\mathcal{M})}.$$

Thus,

$$\|\pi_b(f)\|_{BMO_{cr}(\mathcal{M})}^2 \leq (c_q + 1)(\|\pi_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})} + \|b\|_{BMO_r(\mathcal{M})}) \|\pi_b(f)\|_{BMO_{cr}(\mathcal{M})},$$

Therefore

$$\|\pi_b\|_{L^\infty(\mathcal{M}) \rightarrow BMO_{cr}(\mathcal{M})} \leq (c_q + 1)(\|\pi_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})} + \|b\|_{BMO_r(\mathcal{M})}). \quad \blacksquare$$

*Proof of Theorem 1.2.* By Lemma 3.7 and Lemma 3.8 we get immediately that

$$\begin{aligned} & \max \{ \|\pi_b\|_{L^\infty(\mathcal{M}) \rightarrow BMO_{cr}(\mathcal{M})}, \|\tilde{\pi}_b\|_{L^\infty(\mathcal{M}) \rightarrow BMO_{cr}(\mathcal{M})} \} \\ & \leq c_p \max \{ \|\pi_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})}, \|\tilde{\pi}_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})} \} \end{aligned}$$

By the interpolation results of noncommutative martingales (Theorem 3.5), we get

$$\begin{aligned} & \max \{ \|\pi_b\|_{L^q(\mathcal{M}) \rightarrow L^q(\mathcal{M})}, \|\tilde{\pi}_b\|_{L^q(\mathcal{M}) \rightarrow L^q(\mathcal{M})} \} \\ & \leq c_p \max \{ \|\pi_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})}, \|\tilde{\pi}_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})} \}, \end{aligned}$$

for all  $1 < p < q < \infty$ .

**Question :** Assume  $\pi_b, \tilde{\pi}_b$  are of type  $(p, p)$ , are they of weak type  $(1, 1)$ ? More precisely, assume  $\|\pi_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})} + \|\tilde{\pi}_b\|_{L^p(\mathcal{M}) \rightarrow L^p(\mathcal{M})} < \infty$ , does there exist a constant  $C > 0$  such that, for any  $f \in L^1(\mathcal{M})$ ,  $\lambda > 0$ , there is a projection  $e \in \mathcal{M}$  such that

$$\tau(e^\perp) \leq C \frac{\|f\|_{L^1(\mathcal{M})}}{\lambda} \quad \text{and} \quad \|e\pi_b(f)e\|_{L^\infty(\mathcal{M})} + \|e\tilde{\pi}_b(f)e\|_{L^\infty(\mathcal{M})} \leq \lambda?$$

We have the following corollary by applying results of this section to matrix valued dyadic paraproducts discussed in Section 1 and Section 2. Note  $M_n$  valued dyadic martingales on the unit circle are noncommutative martingales associated with the von Neuman algebra  $\mathcal{M} = L^\infty(\mathbb{T}) \otimes M_n$  and the filtration  $\mathcal{M}_k = L^\infty(\mathbb{T}, \mathcal{F}_k) \otimes M_n$ .

**Corollary 3.9** *Let  $1 < p < \infty$ , denote by  $c_p(n)$  the best constant such that*

$$\|\pi_b\|_{L^p(S_n^p) \rightarrow L^p(S_n^p)} \leq c_p(n) \|b\|_{L^\infty(M_n)}, \quad \forall b.$$

*Then*

$$c_p(n) \asymp \log(n+1).$$

**Proof.** Note in the proof of Theorem 1.1, if we see  $f$  as a column matrix valued function and  $g$  as a row matrix valued function, we will have

$$\|f\|_{L^p(S_n^p)} = \|\alpha\|_{\ell_n^2}, \quad \|g\|_{L^q(S_n^q)} = \|\beta\|_{\ell_n^2}.$$

By the same method, we can prove  $c_p(n) \geq c \log(n+1)$  for all  $1 < p < \infty$ . For the inverse relation, by (1.2) we have  $c_2(n) \leq c \log(n+1)$ . Then, by (3.15), we get

$$\begin{aligned} \|\pi_b\|_{L^\infty(M_n) \rightarrow BMO_{cr}} &\leq c_2(c_2(n) \|b\|_{L^\infty(M_n)} + \|b\|_{BMO_{cr}}) \\ &\leq c \log(n+1) \|b\|_{L^\infty(M_n)}, \quad \forall b \in L^\infty(M_n) \end{aligned} \quad (3.18)$$

Denote by  $\pi_b^*$  the adjoint operator of the dyadic paraproduct  $\pi_b$ , then

$$\pi_b^*(f) = \sum_k (d_k b)^* E_{k-1} f.$$

Note we have the decomposition

$$\pi_b^*(f) = b^* f - \pi_{b^*}(f) - (\pi_{f^*}(b))^*.$$

By (3.18), we get

$$\begin{aligned} \|\pi_b^*\|_{L^\infty(M_n) \rightarrow BMO_{cr}} &\leq \|b^*\|_{L^\infty(M_n)} + c \log(n+1) \|b^*\|_{L^\infty(M_n)} + c \log(n+1) \|b\|_{L^\infty(M_n)} \\ &\leq c \log(n+1) \|b\|_{L^\infty(M_n)}. \end{aligned} \quad (3.19)$$

By (3.18), (3.19) and the interpolation result Theorem 3.5, we get

$$\|\pi_b\|_{L^p(S_n^p) \rightarrow L^p(S_n^p)} \leq c_p \log(n+1) \|b\|_{L^\infty(M_n)}, \quad \forall 1 < p < \infty.$$

Therefore, we can conclude  $c_p(n) \asymp \log(n+1)$ . ■

**Acknowledgments.** The author is indebted to his adviser G. Pisier for many helpful conversations. The example in the proof of Theorem 1.1 comes out from the discussions with him. The author also would like to thank the referee for his very useful comments.

## References

- [1] O. Blasco, S. Pott, Operator-valued dyadic BMO Spaces. Preprint.
- [2] T. A. Gillespie, S. Pott, S. Treil, A. Volberg, Logarithmic growth for martingale transform, *Journal of London Mathematical Society*(2), 64 (2001), no. 3, 624-636.
- [3] M. Junge, M. Musat, Non-commutative John-Nirenberg theorem, *Transaction of AMS*, to appear.
- [4] Nets H. Katz, Matrix valued paraproducts, *J. Fourier Anal. Appl.* 300 (1997), 913–921.
- [5] S. Kwapien, A. Pelczyński, The main triangle projection in matrix spaces and its applications, *Studia Math.* 34 (1970) 43–67.

- [6] M. Musat, Interpolation between noncommutative BMO and noncommutative  $L_p$ -spaces, *Journal of Functional Analysis*, 202 (2003), no. 1, 195-225.
- [7] F. Nazarov, G. Pisier, S. Treil, A. Volberg, Sharp estimates in vector Carleson imbedding theorem and for vector paraproducts. *J. Reine Angew. Math.* 542 (2002), 147-171.
- [8] F. Nazarov, S. Treil, and A. Volberg, Counterexample to infinite dimensional Carleson embedding theorem, *C. R. A. Sci. Paris Ser. I Math.* 325 (1997), no. 4, 383–388.
- [9] St. Petermichl, Dyadic shift and a logarithmic estimate for Hankel operator with matrix symbol, *Compt. Rend. Acad. Sci. Paris* 330 (2000), no. 1, 455–460.
- [10] G. Pisier, Notes on Banach space valued  $H^p$ -spaces, preprint.
- [11] S. Pott, M. Smith, Vector paraproducts and Hankel operators of Schatten class via p-John-Nirenberg theorem, *J. Funct. Anal.* 217(2004), no. 1, 38-78.

Dept. of Math.  
 Texas A&M University  
 College Station, TX, 77843  
 U. S. A.  
 tmei@math.tamu.edu